

# Catalan Numbers

CS 491 – Competitive Programming

**Dr. Mattox Beckman**

UNIVERSITY OF ILLINOIS AT URBANA-CHAMPAIGN  
DEPARTMENT OF COMPUTER SCIENCE

Spring 2024

# Objectives

- ▶ Compute the  $n$ th Catalan number
- ▶ Map the Catalan numbers to various isomorphisms.


## Example 1: Parenthesis

- ▶ Suppose you have a  $2n$  parenthesis characters, half open, half closed. How many “algebraically legal” sequences are there?
  - ▶ Every open parenthesis followed by a matching close parenthesis.
  - ▶ No close parenthesis before its corresponding open.
- ▶  $n = 2$ 
  - ▶  $()()$
  - ▶  $(( ))$
- ▶  $n = 3$ 
  - ▶  $(( ( )))$
  - ▶  $() ( ( ) )$
  - ▶  $() () ()$
  - ▶  $(( )) ()$
  - ▶  $( ( ) ) ()$
- ▶ For  $n = 4$  there are 14.


## Example 2: Trees

- ▶ Given  $n$  indistinguishable nodes, how many binary trees can you make?

- ▶  $n = 2$



- ▶  $n = 3$






## Example 3: Integer Sequences of +1 and -1

- ▶ Suppose you have  $n$  copies of 1 and  $n$  copies of  $-$ .
- ▶ How many sequences of  $a_0, a_1, \dots, a_{2n}$  are there such that for all  $k < 2n$ ,  $0 \leq \sum_{i=0}^k a_i$ ?
- ▶  $n = 2$ 
  - ▶ 1,1,-1,-1
  - ▶ 1,-1,1,-1
- ▶  $n = 3$ 
  - ▶ 1,-1,1,-1,1,-1
  - ▶ 1,-1,1,1,-1,-1
  - ▶ 1,1,-1,-1,1,-1
  - ▶ 1,1,-1,1,-1,-1
  - ▶ 1,1,1,-1,-1,-1


## Example 4: Matrix Walk

- ▶ Suppose you have a  $n \times n$  matrix. Start from the top left and reach the bottom right without crossing the diagonal.

- ▶  $n = 2$



- ▶  $n = 3$



# The Formula

- ▶ Recursively:  $Can(n) = \frac{(2n-1)(2n)}{(n+1)n} Cat(n-1)$ 
  - ▶  $Cat(0) = 1, Cat(1) = 1, Cat(2) = 2, Cat(3) = 5, Cat(4) = 14, \dots$
- ▶ Others things:
  - ▶ number of ways to triangulate a polygon
  - ▶ Number of ways to count a tie vote so that candidate A never passes candidate B